



Volume:01/Issue:10/October-2026

www.paperly-publications.com

RECORD SHEET

PAPERLY ID

PLY-2026-000017

EARLIER ID

RP-2026-000017

DOI

<https://doi.org/10.5281/zenodo.23363308>

VERSION

1.0 · VER-2026-000005

PUBLISHED

14 October 2026

STATUS

Published

RECORD TYPE

Repository record

REVIEW

Peer reviewed (completed reviews recorded)

LICENCE

CC-BY

FILE SHA-256 (ORIGINAL FILE)

f3c3c31a42a3bdce96d4aa742c12cf2de4bb9a9085ce09c0995827667f0d6609

ANATOMY OF THE EYEBALL

1. Raiymova Klara Amangeldievna
2. KARUNAKARAN BAIJU KOUSHIK
3. Rohit Tanwar
4. Anjum Saniya
5. Arumugam Prabanchan

HOW TO CITE

Amangeldievna 1RK, KOUSHIK 2KB, Tanwar 3R, Saniya 4A, Prabanchan 5A.
ANATOMY OF THE EYEBALL [repository record]. Paperly; 2026.
[doi:10.5281/zenodo.23363308](https://doi.org/10.5281/zenodo.23363308)



Scan or open www.paperly-publications.com/article/PLY-2026-000017 to see the record, its version history and any notices.

The manuscript was submitted as a Word file. The pages after this sheet are a text rendering of it (images and complex layout are left out).

This sheet and the margin stamp were added by Paperly. They are not part of the manuscript. The Paperly ID is an internal identifier, not a DOI. A certificate ID can be checked at the verification address behind it.

Prepared 14 October 2026.

ANATOMY OF THE EYEBALL

- Raiymova Klara Amangeldievna
- KARUNAKARAN BAIJU KOUSHIK
- Rohit Tanwar
- Anjum Saniya
- Arumugam Prabanchan

1.LECTURER, INTERNATIONAL MEDICAL FACULTY, OSH STATE UNIVERSITY, OSH KYRGYZ REPUBLIC.

2,3,4,5, STUDENTS, INTERNATIONAL MEDICAL FACULTY, OSH STATE UNIVERSITY, OSH KYRGYZ REPUBLIC.

1. ABSTRACT

The eyeball is a highly specialized sensory organ responsible for the perception of light and the formation of visual images. Its complex anatomical organization enables the conversion of external light stimuli into electrical signals that are transmitted to the brain through the optic nerve. A comprehensive understanding of the anatomy of the eyeball is essential for medical students and healthcare professionals because structural abnormalities of the eye are associated with several ophthalmological and systemic disorders. The aim of this review is to describe the gross anatomy, microscopic structure, anatomical relationships, blood supply, nerve supply, and clinical significance of the human eyeball. A narrative literature review was conducted using standard anatomical and ophthalmological textbooks, peer-reviewed scientific publications, and established medical references. The anatomical findings were organized according to the three principal coats of the eyeball: the fibrous coat, vascular coat (uvea), and nervous coat (retina). The sclera maintains structural integrity, while the choroid provides vascular support to the outer retina. The anatomical organization of the anterior and posterior chambers, vitreous body, and optic nerve is also essential for normal visual function. A thorough understanding of these structures provides the anatomical foundation for diagnosing and managing conditions such as glaucoma, cataract, retinal detachment, and optic neuropathy.

2. KEYWORDS

- Eyeball
- Eye Anatomy
- Retina
- Cornea
- Optic Nerve
- Ocular Structures

3. INTRODUCTION

The human eye is a specialized sensory organ responsible for vision. It detects light, distinguishes colours, recognizes shapes, perceives depth, and provides information about the surrounding environment. Vision is essential for daily activities, communication, education, and

overall quality of life.

Anatomically, the eyeball (bulbus oculi) is a nearly spherical structure situated within the bony orbit. It is surrounded by protective structures, including the eyelids, conjunctiva, orbital fat, extraocular muscles, and lacrimal apparatus. The adult eyeball measures approximately 24 mm in the anteroposterior diameter, although its dimensions vary among individuals. Its weight is approximately 7–8 g. The eyeball is connected to the brain through the optic nerve, which transmits visual information to the central nervous system [1,2].

The eye is anatomically divided into three principal coats:

- Fibrous coat: The outer protective layer, consisting of the sclera and cornea.
- Vascular coat (uvea): The middle layer, consisting of the choroid, ciliary body, and iris.
- Nervous coat: The innermost layer, represented by the retina.

Within these coats are specialized structures, including the crystalline lens, aqueous humour, vitreous body, and optic nerve head. These structures work together to maintain the optical properties and physiological functions of the eye [2,3].

The eyeball is located in the anterior part of the orbit, with approximately one-sixth of its anterior surface exposed and the remaining five-sixths protected by the orbital margins and surrounding tissues. The anterior pole is the central point of the corneal curvature, while the posterior pole is the corresponding point on the posterior surface of the eyeball. The equator is an imaginary circular line midway between the anterior and posterior poles [1,4].

Global and Regional Relevance

Visual impairment represents an important global public health concern. According to the World Health Organization, at least 2.2 billion people worldwide have near or distance vision impairment, and in at least 1 billion cases, the impairment could have been prevented or remains unaddressed [5].

Major causes of visual impairment include refractive errors, cataracts, glaucoma, diabetic retinopathy, age-related macular degeneration, and corneal diseases [5,6].

A sound knowledge of ocular anatomy is particularly important in regions where access to specialized ophthalmological services may be limited. Understanding the anatomical basis of eye diseases helps medical practitioners recognize early clinical manifestations, perform appropriate examinations, and refer patients for timely treatment.

Rationale of the Review

The eyeball contains several interconnected anatomical structures, each performing specific functions. Abnormalities involving even a small component may significantly affect vision. For example, corneal opacity interferes with light transmission, lens opacity produces cataract, and damage to retinal photoreceptors may cause permanent visual impairment.

4. METHODOLOGY

4.1 Study Design

This study was designed as a descriptive narrative literature review of the anatomy of the human eyeball. It involved the examination and synthesis of established anatomical and ophthalmological literature rather than an original clinical or experimental investigation.

4.2 Study Population

No human participants or animal subjects were recruited. The subject of the review was the normal anatomical structure of the adult human eyeball.

4.3 Inclusion Criteria

The following sources were considered:

- Standard medical anatomy textbooks.
- Established ophthalmology textbooks.
- Peer-reviewed publications concerning ocular anatomy and physiology.
- Scientific literature describing the structure and function of the cornea, sclera, uvea, retina, lens, and optic nerve.
- Authoritative publications concerning visual impairment and ocular health.

4.4 Exclusion Criteria

The following materials were excluded:

- Unverified internet sources.
- Non-scientific articles and unsupported anatomical descriptions.
- Publications containing insufficient anatomical information.
- Studies primarily concerned with unrelated systemic diseases.
- Experimental findings that could not be appropriately applied to normal human ocular anatomy.

4.5 Data Collection Methods

Relevant anatomical information was obtained from standard textbooks, including Gray's Anatomy, Snell's Clinical Anatomy, and established ophthalmology references.

The information was classified into the following categories:

- Gross anatomy of the eyeball.
- Fibrous coat.
- Vascular coat.
- Nervous coat.
- Lens and refractive structures.
- Ocular chambers and fluids.
- Blood supply and innervation.
- Clinical and anatomical correlations.

4.6 Statistical Analysis

No statistical analysis was performed because the review did not involve primary quantitative data collection. Anatomical descriptions were compared and organized according to established medical literature.

4.7 Ethical Considerations

The study involved only previously published scientific and educational literature. No patient information, personal data, or biological specimens were used. Therefore, institutional ethical approval and informed consent were not required.

5. RESULTS

The literature review identified a consistent anatomical organization of the human eyeball into three principal coats, together with specialized internal structures responsible for optical refraction, accommodation, and visual signal transmission.

5.1 General Anatomical Characteristics

The adult human eyeball is approximately spherical and measures around 24 mm in its anteroposterior diameter. It is situated within the orbit and is connected posteriorly to the brain through the optic nerve.

Table 1. General anatomical characteristics of the eyeball

Anatomical parameter	Description
Shape	Nearly spherical
Anteroposterior diameter	Approximately 24 mm
Weight	Approximately 7–8 g
Location	Bony orbit
Anterior pole	Central point of corneal curvature
Posterior pole	Corresponding posterior point
Equator	Imaginary circumference midway between the poles
Principal coats	Fibrous, vascular, and nervous
Main sensory function	Vision

5.2 Three Coats of the Eyeball

Table 2. Anatomical organization of the three coats

Coat	Components	Principal function
Fibrous coat	Sclera and cornea	Protection, shape, and refraction
Vascular coat	Choroid, ciliary body, and iris	Nutrition, accommodation, and regulation of light
Nervous coat	Retina	Reception of light and conversion into neural signals

5.3 Internal Structures

The internal structures of the eyeball include the lens, aqueous humour, vitreous body, and optic disc. These structures are responsible for maintaining optical transparency, focusing light, maintaining intraocular pressure, and transmitting visual information.

Table 3. Internal structures and their functions

Structure | Anatomical location | Principal function

Lens | Behind the iris | Focuses light on the retina

Aqueous humour | Anterior and posterior chambers | Maintains intraocular pressure and nourishes avascular tissues

Vitreous body | Posterior segment | Maintains globe shape and supports the retina

Optic disc | Posterior retina | Exit of optic nerve fibres

Macula lutea | Central posterior retina | Central vision

Fovea centralis | Centre of macula | Highest visual acuity

5.4 Measurements of Important Ocular Structures

Table 4. Approximate anatomical measurements

Structure | Approximate measurement

Adult eyeball diameter | 24 mm

Corneal diameter | 11.5–12 mm horizontally

Central corneal thickness | Approximately 520–550 μm

Lens thickness | Approximately 4 mm in a young adult

Lens diameter | Approximately 9–10 mm

Central anterior chamber depth | Approximately 3 mm

Optic nerve length within orbit | Approximately 25–30 mm

These values are approximate and vary with age, sex, refractive status, and individual anatomical characteristics.

5.5 Blood Supply

The principal arterial supply of the eyeball is derived from the ophthalmic artery, a branch of the internal carotid artery. Its branches include the central retinal artery and posterior ciliary arteries.

The central retinal artery supplies the inner retinal layers, while the choroidal circulation supplies the outer retina, including the photoreceptor region.

5.6 Nerve Supply

The sensory innervation of the eyeball is mainly provided by the ophthalmic division of the trigeminal nerve (CN V1). The autonomic nervous system regulates pupillary function, ciliary muscle activity, and vascular responses.

The optic nerve (CN II) carries visual information from the retina to the brain.

6. DESCRIPTION / DISCUSSION

6.1 Gross Anatomy of the Eyeball

The eyeball is a specialized sensory organ enclosed within the bony orbit. Its approximately spherical shape allows coordinated optical function and movement.

The anterior surface is covered by the transparent cornea, which is continuous at the limbus with the opaque sclera. The posterior aspect is penetrated by the optic nerve.

The eyeball is divided into anterior and posterior segments. The anterior segment contains the cornea, anterior chamber, iris, posterior chamber, and lens. The posterior segment contains the vitreous body, retina, choroid, and posterior part of the sclera [1,2].

Anatomically, the following landmarks are important:

Anterior pole: The central point of the anterior corneal curvature.

Posterior pole: The corresponding point on the posterior surface of the globe.

Equator: An imaginary circular line that divides the eyeball into anterior and posterior halves.

Visual axis: The line connecting the fixation point with the fovea through the nodal region of the optical system.

Optic axis: An idealized line passing through the centres of curvature of the optical surfaces.

These anatomical and optical axes are not identical [2,3].

6.2 Fibrous Coat of the Eyeball

The fibrous coat is the outermost layer of the eyeball. It consists of the sclera and cornea. Its main functions are to protect internal structures, maintain the shape of the globe, and provide attachment to the extraocular muscles [1,4].

A. Sclera

The sclera is the white, opaque, tough connective tissue forming approximately the posterior five-sixths of the fibrous coat.

It is composed mainly of dense irregular collagenous connective tissue, with fibroblasts and elastic fibres.

Anatomical features:

- It is relatively thick posteriorly and becomes thinner near the equator.
- It is continuous anteriorly with the cornea at the corneoscleral junction.
- Posteriorly, it is perforated by the optic nerve.
- It provides attachment to the six extraocular muscles.
- Its external surface is covered by the episclera and Tenon's capsule.

Layers of the sclera:

- Episclera: The superficial vascular connective tissue layer.
- Scleral stroma: The thick, dense collagenous middle layer.
- Lamina fusca: The inner pigmented layer adjacent to the choroid.

The posterior sclera contains a perforated region called the lamina cribrosa, through which retinal ganglion cell axons pass to form the optic nerve [1,4].

Clinical significance:

In glaucoma, increased intraocular pressure and other biomechanical factors can produce structural changes in the optic nerve head and lamina cribrosa. Scleral inflammation is known as scleritis and may be associated with autoimmune diseases [7,8].

B. Cornea

The cornea is the transparent anterior one-sixth of the fibrous coat. It is avascular and receives nutrients from the aqueous humour, tears, and peripheral limbal circulation.

Its major function is to refract incoming light. The cornea contributes approximately two-thirds of the total refractive power of the eye, with a typical power of approximately +43 dioptres [3,4].

Layers of the cornea:

The classical anatomical description identifies five layers:

- Corneal epithelium.
- Bowman's layer.
- Corneal stroma.
- Descemet's membrane.
- Corneal endothelium.

Corneal epithelium:

It is a non-keratinized stratified squamous epithelium consisting of approximately five to seven cell layers. It provides a protective barrier against microorganisms and mechanical injury.

Bowman's layer:

This is an acellular layer located beneath the epithelium. It consists of densely arranged collagen fibres.

Corneal stroma:

The stroma forms approximately 90% of the corneal thickness. It contains regularly arranged collagen lamellae, keratocytes, and extracellular matrix. Its organized structure is essential for transparency.

Descemet's membrane:

It is the basement membrane of the corneal endothelium.

Corneal endothelium:

It is a single layer of polygonal cells that maintains corneal dehydration through barrier and pump functions. Damage to endothelial cells can cause corneal oedema and loss of transparency [3,4].

Clinical significance:

Corneal ulceration, keratitis, trauma, and endothelial dysfunction can produce corneal opacity and impaired vision. Corneal transplantation may be required in selected cases of severe corneal damage.

6.3 Vascular Coat (Uveal Tract)

The vascular coat is the middle layer of the eyeball and is also known as the uvea. It consists of three parts:

- Choroid.
- Ciliary body.
- Iris.

The uvea is highly vascular and contributes to ocular nutrition, accommodation, and regulation of the amount of light entering the eye [1,3].

A. Choroid

The choroid is a highly vascular, pigmented connective tissue layer situated between the sclera and retina.

It extends from the optic nerve posteriorly to the ciliary body anteriorly.

Functions:

- Supplies oxygen and nutrients to the outer retinal layers.
- Removes metabolic waste products.
- Absorbs scattered light through its melanin content.
- Helps maintain the metabolic environment of the photoreceptors.

Layers of the choroid:

- Suprachoroidal layer.
- Haller's layer.
- Sattler's layer.
- Choriocapillaris.
- Bruch's membrane, which forms the interface between the choroid and retinal pigment epithelium.

The choriocapillaris consists of fenestrated capillaries and plays a major role in supplying the outer retina [3,4].

Clinical significance:

Inflammation of the choroid is called choroiditis. Involvement of the choroid and retina is termed chorioretinitis. Choroidal vascular abnormalities may also contribute to retinal disorders.

B. Ciliary Body

The ciliary body is the anterior continuation of the choroid. It extends from the ora serrata posteriorly to the root of the iris anteriorly.

It is triangular in cross-section and contains smooth muscle and vascular connective tissue.

The ciliary body is divided into two parts:

- Pars plicata: The anterior folded region containing ciliary processes.
- Pars plana: The relatively flat posterior region.

Components:

The ciliary body consists of the ciliary muscle, stroma, and ciliary epithelium.

Functions of the ciliary body:

- Produces aqueous humour through the ciliary processes.
- Participates in accommodation through contraction and relaxation of the ciliary muscle.
- Provides attachment to the suspensory ligaments of the lens.

Accommodation:

During near vision, the parasympathetic nervous system stimulates contraction of the ciliary muscle. This reduces tension on the zonular fibres, allowing the lens to become more convex and increase its refractive power.

During distance vision, the ciliary muscle relaxes, zonular tension increases, and the lens becomes flatter [2,3].

Clinical significance:

Age-related reduction in lens elasticity produces presbyopia. Dysfunction of the ciliary body can affect aqueous humour production and accommodation.

C. Iris

The iris is the coloured, circular diaphragm located behind the cornea and in front of the lens.

It contains a central opening called the pupil, through which light enters the eye.

The iris separates the anterior and posterior chambers of the anterior segment.

Components of the iris:

- Anterior border layer.
- Iris stroma.
- Sphincter pupillae muscle.
- Dilator pupillae muscle.
- Posterior pigmented epithelium.

Muscles of the iris:

- Sphincter pupillae: A circular smooth muscle that constricts the pupil.
- Dilator pupillae: A radially arranged smooth muscle that dilates the pupil.

Nerve supply:

- Parasympathetic fibres cause pupillary constriction through the oculomotor nerve.
- Sympathetic fibres cause pupillary dilation through the superior cervical ganglion pathway.

Clinical significance:

Abnormal pupillary size or reaction may indicate neurological disorders, pharmacological effects, or ocular disease. Examination of pupillary light reflexes is an important component of neurological assessment [2,7].

6.4 Nervous Coat: Retina

The retina is the innermost nervous layer of the eyeball. It is responsible for detecting light and converting it into electrical signals.

It extends from the optic disc posteriorly to the ora serrata anteriorly. The visual part of the retina contains photoreceptors, whereas the anterior non-visual part continues over the ciliary body and posterior surface of the iris [3,4].

The retina consists of two major components:

- Retinal pigment epithelium.
- Neural retina.

Layers of the Retina

The retina is classically described as having ten histological layers, from outside to inside:

- Retinal pigment epithelium.
- Layer of rods and cones.
- External limiting membrane.
- Outer nuclear layer.
- Outer plexiform layer.
- Inner nuclear layer.
- Inner plexiform layer.
- Ganglion cell layer.
- Nerve fibre layer.
- Internal limiting membrane.

Retinal pigment epithelium:

It is a single layer of pigmented epithelial cells located between the neural retina and choroid. It supports photoreceptor function, participates in the visual cycle, absorbs excess light, and phagocytoses shed photoreceptor outer segments.

Photoreceptors:

There are two principal types:

- Rods: Responsible mainly for dim-light vision and peripheral visual sensitivity.
- Cones: Responsible for colour vision and high-resolution vision under brighter illumination.

Bipolar cells:

These neurons transmit signals from photoreceptors to ganglion cells.

Ganglion cells:

They receive signals from retinal interneurons. Their axons form the nerve fibre layer and converge at the optic disc to form the optic nerve [3,4].

Important Anatomical Regions of the Retina

A. Optic Disc

The optic disc is the region where retinal ganglion cell axons leave the eyeball to form the optic nerve.

It is also the site of entry and exit of the central retinal vessels.

The optic disc lacks rods and cones, producing the physiological blind spot.

B. Macula Lutea

The macula lutea is a specialized central retinal region responsible for detailed central vision and colour discrimination.

It contains a high concentration of cones.

C. Fovea Centralis

The fovea centralis is a small depression situated in the centre of the macula. It contains densely packed cones and is responsible for the highest visual acuity.

D. Ora Serrata

The ora serrata is the serrated anterior boundary of the photosensitive retina. It marks the transition between the neural retina and the non-visual ciliary part of the retina.

Clinical significance:

Retinal detachment occurs when the neurosensory retina separates from the retinal pigment epithelium. Diabetic retinopathy is associated with retinal microvascular damage, while age-related macular degeneration affects the macular region [6,8].

6.5 Lens of the Eye

The crystalline lens is a transparent, biconvex, avascular structure situated behind the iris and in front of the vitreous body.

It is suspended by zonular fibres attached to the ciliary body.

The lens is composed of:

- Lens capsule.
- Anterior lens epithelium.
- Lens fibres.

The lens capsule is a transparent basement membrane surrounding the lens. The anterior surface contains a single layer of epithelial cells, which differentiate into lens fibres.

The lens is responsible for fine adjustment of the refractive power of the eye during accommodation [2,3].

Clinical significance:

Cataract refers to opacity of the crystalline lens. It can cause progressive visual impairment, glare, and reduced contrast sensitivity. Surgical removal of the opaque lens with intraocular lens implantation is the standard treatment for visually significant cataract [6,9].

6.6 Chambers of the Eyeball

The internal spaces of the anterior segment contain aqueous humour, while the posterior segment is largely occupied by vitreous humour.

The eyeball is divided into three principal spaces:

A. Anterior Chamber

The anterior chamber lies between the posterior surface of the cornea and the anterior surface of the iris.

It contains aqueous humour.

B. Posterior Chamber

The posterior chamber is a narrow space situated behind the iris and in front of the lens and zonular apparatus.

It communicates with the anterior chamber through the pupil.

C. Vitreous Chamber

The vitreous chamber is the largest internal compartment. It is located behind the lens and is surrounded by the retina, ciliary body, and posterior lens surface.

It contains the vitreous body [2,3].

6.7 Aqueous Humour

Aqueous humour is a transparent fluid produced mainly by the non-pigmented ciliary epithelium of the ciliary processes.

Functions:

- Maintains intraocular pressure.
- Provides nutrients to the avascular cornea and lens.
- Removes metabolic waste products.
- Maintains the optical environment of the anterior segment.

Formation and circulation:

Aqueous humour is secreted into the posterior chamber. It flows through the pupil into the anterior chamber and drains mainly through the trabecular meshwork into Schlemm's canal and the episcleral venous system.

A smaller proportion drains through the uveoscleral pathway [3,7].

Clinical significance:

Increased resistance to aqueous humour drainage can elevate intraocular pressure and contribute to glaucoma. Glaucoma is characterized by progressive optic neuropathy and may lead to irreversible visual field loss.

6.8 Vitreous Body

The vitreous body is a transparent, gelatinous substance occupying the vitreous chamber.

It consists mainly of water, with collagen fibres, hyaluronic acid, and other extracellular matrix components.

Functions:

- Maintains the shape of the eyeball.
- Provides a transparent medium for light transmission.
- Helps support the retina.

- Participates in the internal structural organization of the posterior segment.

With increasing age, the vitreous undergoes liquefaction and structural changes. Posterior vitreous detachment may occur, sometimes producing flashes of light and floaters [3,8].

6.9 Blood Supply of the Eyeball

The blood supply of the eyeball is primarily derived from the ophthalmic artery, which arises from the internal carotid artery.

The principal arterial branches include:

1. Central Retinal Artery

The central retinal artery enters the optic nerve and reaches the retina through the optic disc. It supplies the inner retinal layers.

It is functionally an end artery, and its occlusion can cause sudden painless loss of vision.

2. Short Posterior Ciliary Arteries

These arteries supply the choroid, optic nerve head, and adjacent structures.

3. Long Posterior Ciliary Arteries

These arteries contribute to the blood supply of the anterior uveal tract and participate in the formation of the major arterial circle of the iris.

4. Anterior Ciliary Arteries

These vessels arise from muscular branches of the ophthalmic circulation and supply the anterior segment and contribute to the arterial circulation of the iris and ciliary body.

The retinal and choroidal circulations have distinct anatomical arrangements [1,3].

Venous drainage:

The principal venous drainage occurs through:

- Central retinal vein.
- Vortex veins.
- Anterior ciliary venous channels.

The vortex veins drain the choroidal circulation into the ophthalmic venous system [1,4].

6.10 Nerve Supply of the Eyeball

The nerve supply of the eyeball includes sensory, parasympathetic, and sympathetic components.

A. Sensory Nerve Supply

The sensory supply is mainly derived from the ophthalmic division of the trigeminal nerve (CN V1).

The nasociliary nerve gives rise to the long ciliary nerves and contributes sensory fibres to the eyeball.

The long and short ciliary nerves provide sensory innervation to the cornea and other ocular structures.

B. Parasympathetic Nerve Supply

Parasympathetic fibres originate from the Edinger–Westphal nucleus and travel with the oculomotor nerve (CN III).

They synapse in the ciliary ganglion.

Postganglionic fibres travel through the short ciliary nerves and supply:

- Sphincter pupillae.
- Ciliary muscle.

Their functions include pupillary constriction and accommodation.

C. Sympathetic Nerve Supply

Sympathetic fibres originate from the hypothalamus and descend to the ciliospinal centre. They synapse in the superior cervical ganglion.

Postganglionic fibres reach the eyeball through the internal carotid plexus and ciliary nerves.

They supply the dilator pupillae and contribute to ocular vascular regulation [2,4].

6.11 Optic Nerve and Visual Pathway

The optic nerve is the second cranial nerve and carries visual information from the retina to the brain.

It is formed by the axons of retinal ganglion cells.

The optic nerve is approximately 40–50 mm long from the globe to the optic chiasm, although the measured length depends on the anatomical segment included.

It is divided into four portions:

- Intraocular portion.
- Intraorbital portion.
- Intracanalicular portion.
- Intracranial portion.

At the optic chiasm, fibres originating from the nasal halves of both retinas cross to the opposite side, whereas temporal retinal fibres remain uncrossed.

The optic tracts carry visual information to the lateral geniculate bodies of the thalamus. From there, optic radiations project to the primary visual cortex in the occipital lobe [2,3].

Clinical significance:

Damage to different parts of the visual pathway produces characteristic visual field defects. For example, optic chiasm compression may produce bitemporal hemianopia, while lesions of the optic tract may produce contralateral homonymous hemianopia.

6.12 Clinical Importance of Eyeball Anatomy

Understanding ocular anatomy is essential for recognizing common ophthalmological disorders.

Table 5. Anatomical structures and associated clinical conditions

Anatomical structure | Clinical condition | Anatomical basis

Cornea | Keratitis | Inflammation of corneal tissue

Cornea | Corneal opacity | Loss of corneal transparency

Sclera | Scleritis | Inflammation of scleral tissue

Lens | Cataract | Opacity of the crystalline lens

Iris | Iritis | Inflammation of the iris

Ciliary body | Cyclitis | Inflammation of the ciliary body

Trabecular meshwork | Glaucoma | Increased resistance to aqueous humour drainage in many forms

Retina | Retinal detachment | Separation of neurosensory retina from retinal pigment epithelium

Macula | Macular degeneration | Damage involving the macular region

Optic nerve | Optic neuropathy | Damage to optic nerve fibres

7.1 Clinical Recommendations

- Routine eye examinations should include visual acuity assessment, pupillary reactions, and examination of the anterior segment.
- Fundoscopic examination should be performed when clinically indicated to assess the retina and optic disc.
- Patients with diabetes mellitus and other risk factors should undergo appropriate retinal screening.
- Sudden visual loss, new-onset flashes and floaters, or a curtain-like visual field defect require urgent ophthalmological assessment.
- Patients with suspected glaucoma should undergo appropriate optic nerve and intraocular pressure evaluation.

7.2 Preventive Recommendations

- Promote awareness regarding eye injuries and the use of appropriate protective eyewear.
- Encourage regular ophthalmological examinations in individuals at increased risk of ocular disease.
- Promote diabetes and hypertension control to reduce the risk of associated ocular complications.
- Encourage early diagnosis and management of refractive errors and other correctable visual disorders.
- Improve access to ophthalmological services, particularly in underserved populations.

7.4 Recommendations for Future Research

- Further anatomical studies should examine age-related variations in ocular structures.
- Advanced imaging techniques, including optical coherence tomography, should be used to study retinal anatomy.

- Research should investigate the relationship between ocular anatomical variations and glaucoma.
- More population-based research is needed to understand regional patterns of preventable visual impairment.
- Anatomical teaching models and digital learning resources should be evaluated for their effectiveness in undergraduate medical education.

8. CONCLUSION

The human eyeball is a complex sensory organ with a highly specialized anatomical organization. It consists of three principal coats: the fibrous coat, vascular coat, and nervous coat. The cornea and lens provide optical refraction, the sclera maintains structural integrity, the uvea supplies nutrients and regulates accommodation and pupil size, and the retina converts light into neural signals.

The aqueous humour and vitreous body maintain the internal environment and structural organization of the eyeball. The ophthalmic circulation provides blood supply, while the trigeminal, oculomotor, sympathetic, and optic nerve pathways contribute to sensory, autonomic, and visual functions.

A thorough understanding of the anatomy of the eyeball is fundamental for MBBS students because it forms the basis for understanding common ophthalmological conditions, including cataract, glaucoma, retinal detachment, corneal diseases, and optic neuropathy.

Therefore, knowledge of normal ocular anatomy is essential for accurate clinical examination, early diagnosis, appropriate referral, and effective management of eye disorders.

10. REFERENCES

- Standring S, editor. Gray's Anatomy: The Anatomical Basis of Clinical Practice. 42nd ed. London: Elsevier; 2020.
- Snell RS. Clinical Anatomy by Regions. 10th ed. Philadelphia: Wolters Kluwer; 2019.
- Remington LA. Clinical Anatomy and Physiology of the Visual System. 3rd ed. St Louis: Elsevier Butterworth-Heinemann; 2012.
- Forrester JV, Dick AD, McMenamin PG, Roberts F, Pearlman E. The Eye: Basic Sciences in Practice. 4th ed. Edinburgh: Elsevier; 2016.
- World Health Organization. World report on vision. Geneva: World Health Organization; 2019.
- Bourne RRA, Flaxman SR, Braithwaite T, Cicinelli MV, Das A, Jonas JB, et al. Magnitude, temporal trends, and projections of the global prevalence of blindness and distance and near vision impairment: a systematic review and meta-analysis. Lancet Glob Health. 2017;5(9).
- American Academy of Ophthalmology. Basic and Clinical Science Course. Section 2: Fundamentals and Principles of Ophthalmology. San Francisco: American Academy of Ophthalmology.
- American Academy of Ophthalmology. Basic and Clinical Science Course. Section 12: Retina and Vitreous. San Francisco: American Academy of Ophthalmology.
- American Academy of Ophthalmology. Basic and Clinical Science Course. Section 11: Lens and Cataract. San Francisco: American Academy of Ophthalmology.

- American Academy of Ophthalmology. Basic and Clinical Science Course. Section 10: Glaucoma. San Francisco: American Academy of Ophthalmology.
- Oyster CW. The Human Eye: Structure and Function. Sunderland (MA): Sinauer Associates; 1999.
- Kanski JJ, Bowling B. Clinical Ophthalmology: A Systematic Approach. 8th ed. Edinburgh: Elsevier; 2016.
- Kolb H, Fernandez E, Nelson R, editors. Webvision: The Organization of the Retina and Visual System. Salt Lake City (UT): University of Utah Health Sciences Center; 1995.
- Levin LA, Nilsson SFE, Ver Hoeve J, Wu SM, Kaufman PL, Alm A, editors. Adler's Physiology of the Eye. 11th ed. Edinburgh: Saunders Elsevier; 2011.
- Netter FH. Atlas of Human Anatomy. 7th ed. Philadelphia: Elsevier; 2019.
- Junqueira LC, Carneiro J. Basic Histology: Text and Atlas. 14th ed. New York: McGraw-Hill Education; 2016.
- Moore KL, Dalley AF, Agur AMR. Clinically Oriented Anatomy. 8th ed. Philadelphia: Wolters Kluwer; 2018.
- Snell RS, Lemp MA. Clinical Anatomy of the Eye. 2nd ed. Malden (MA): Blackwell Science; 1998.